
TITLE: MEANS AND METHODS FOR IDENTIFYING GENES AND PROTEINS INVOLVED IN THE PREVENTION AND/OR REPAIR OF A REPLICATION ERROR

Inventor: Tijsterman et al. Docket No.: 2183-6201US

1/7




TITLE: MEANS AND METHODS FOR IDENTIFYING GENES AND PROTEINS INVOLVED IN THE PREVENTION AND/OR REPAIR

OF A REPLICATION ERROR

Inventor: Tijsterman et al. Docket No.: 2183-6201US


2/7

Fig. 1B



Inventor: Tijsterman et al. Docket No.: 2183-6201US

3/7



msh-6 genetic mutants and wildtype C. elegans exposed to msh-6 dsRNA. Spontaneous germline mutation frequency in wildtype C. elegans,

Fig. 2

BEST AVAILABLE COPY

Inventor: Tijsterman et al. Docket No.: 2183-6201US

4/7




Fig. 3: Outline of the principle to detect somatic repeat instability

TITLE: MEANS AND METHODS FOR IDENTIFYING GENES AND PROTEINS INVOLVED IN THE PREVENTION AND/OR REPAIR OF A REPLICATION ERROR

Inventor: Tijsterman et al. Docket No.: 2183-6201US

5/7




Figure 4: A) Wildtype C. elegans containing the in-frame construct, B) the +1 out of frame construct. C) Genetic msh-6 mutants that contain the +1 out of frame construct display LacZ expression.

BEST AVAILABLE COPY

TITLE: MEANS AND METHODS FOR IDENTIFYING GENES AND PROTEINS INVOLVED IN THE PREVENTION AND/OR REPAIR OF A REPLICATION ERROR

Inventor: Tijsterman et al. Docket No.: 2183-6201US

6/7






Fig 5: C. elegans populations fed on E. coli that produce dsRNA homologues to the C. elegans genes unc-22 (A) and msh-6 (B)

BEST AVAILABLE COPY

Inventor: Tijsterman et al. Docket No.: 2183-6201US

7/7



to identify novel mutator loci: Individual animals are fed on dsRNA producing Fig. 6: Schematic representation of the high throughput RNAi based screens bacteria, the progeny is collected and assayed for beta-galactosidase activity